



Immunization Timing

Age 2 months	Interval from previous dose
DTaP	
Hib	
PCV (Pneumo)	
Polio (IPV)	
RV ² (Rotavirus)	
HepB ¹	

Age 4 months	Interval from previous dose
DTaP	1-2 months
Hib	1-2 months
PCV (Pneumo)	1-2 months
Polio (IPV)	1-2 months
RV ² (Rotavirus)	4-10 weeks
HepB ¹	

Age 6 months	Interval from previous dose
DTaP	1-2 months
Hib ³	1-2 months
PCV (Pneumo)	1-2 months
Polio ⁶ (IPV)	1-14 months
RV ² (Rotavirus)	4-10 weeks
HepB ¹	

Age 12 months	Interval from previous dose
DTaP ⁴	
Hib ⁴	2-9 months
PCV ⁴ (Pneumo)	2-6 months
MMR ⁵	
Varicella ⁵ (Chickenpox)	
HepA	

Age 18 months	Interval from previous dose
HepA	6-12 months

Age 6 mos and older	and high risk kids
Flu	Every Year

Early Childhood Program Entry (Age Appropriate)
DTaP, Polio, Hib, Pneumococcal, Hepatitis A, Hepatitis B, MMR, Varicella

Kindergarten Entry
DTaP, Polio, MMR, Varicella, Hepatitis A, Hepatitis B

7th Grade Entry
Varicella, Tdap booster, Hepatitis A, Hepatitis B



Utah Department of Health
IMMUNIZATION PROGRAM
Immunize for healthy lives

Reproduced with permission from the California Department of Health Services, Immunization Branch.

1-800-275-0659 • www.immunize-utah.org

This is a suggested schedule. Your child's immunization schedule may vary depending upon your health care provider's judgment. For more information, consult with your health care provider and/or the "Recommended Childhood Immunization Schedule, United States, 2009" at www.cdc.gov/vaccines.

- ¹ Hepatitis B series: The Hepatitis B vaccine is a three dose series. If Hepatitis B vaccine is not given at birth, the first dose can be given at 2 months of age. The second dose should be given at 4 months of age. The third dose should be given at ≥ 24 weeks of age. If the combination vaccine ComVax[®] (combined Hib and Hepatitis B) is used, a fourth dose can be given. When using ComVax[®], the final dose is given at 12-15 months of age.
- ² RotaTaq[®] vaccine is a three-dose series given at 2, 4, and 6 months of age. Rotarix[®] vaccine is a two-dose series given at 2 and 4 months of age. Vaccination with either rotavirus vaccine should not be started for infants 15 weeks of age or older. All doses should be completed by 8 months of age.
- ³ The number of doses in the Hib series depends on the type of vaccine used. This six month dose of Hib vaccine can be skipped if certain combination vaccines (PedVax[®] and/or ComVax[®]) are used.
- ⁴ The final dose of DTaP, Hib and Pneumococcal (PCV) vaccines should be given ≥ 12 months of age. For school entry, one dose of DTaP should be given ≥ 4 years of age.
- ⁵ The combination vaccine, MMRV (Measles, Mumps, Rubella, Varicella), can be given when both the MMR and the Varicella (Chickenpox) vaccines are necessary.
- ⁶ If Pentacel[®] (DTaP, IPV, Hib) vaccine is used, a fourth dose of IPV at 12-15 months of age may be given. For school entry, one dose of IPV should be given ≥ 4 years of age.