




Childhood Immunization Timing



1-800-275-0659 | www.immunize-utah.org

See reverse for licensed daycare and school entry requirements.

For more information, consult with your health care provider and/or the “Recommended Childhood Immunization Schedule, United States, 2014” at www.cdc.gov/vaccines.

- 1 Hepatitis B series: The Hepatitis B vaccine is a three-dose series. If Hepatitis B vaccine is not given at birth, the 1st dose can be given at 2 months of age. The second dose should be given at 4 months of age. The third dose should be given at ≥ 24 weeks of age. If the combination vaccine ComVax[®] (combined Hib and Hepatitis B) is used, a fourth dose can be given. When using ComVax[®], the final dose is given at 12-15 months of age.
- 2 RotaTeq[®] vaccine is a three-dose series given at 2, 4, and 6 months of age. Rotarix[®] vaccine is a two-dose series given at 2 and 4 months of age. Vaccination with either rotavirus vaccine should not be started for infants 15 weeks of age or older. All doses should be completed by 8 months of age.
- 3 The number of doses in the Hib series depends on the type of vaccine used. This six month dose of Hib vaccine can be skipped if certain combination vaccines (PedVax[®] and/or ComVax[®]) are used.
- 4 If Pentacel[®] (DTaP, IPV, Hib) vaccine is used, a fourth dose of IPV at 12-15 months of age may be given. For school entry, one dose of IPV should be given ≥ 4 years of age.
- 5 The final dose of Hib and Pneumococcal (PCV) vaccines should be given ≥ 12 months of age.
- 6 The fourth dose of DTaP may be given as early as age 12 months of age, provided at least 6 months have passed since the third dose. For school entry, one dose of DTaP should be given ≥ 4 years of age.

1-800-275-0659 | www.immunize-utah.org

Early Childhood Program Entry

(Age Appropriate)

DTaP | Polio | Hib | Pneumococcal
Hepatitis A | Hepatitis B
MMR | Varicella

Kindergarten Entry

DTaP | Polio | MMR | Varicella
Hepatitis A | Hepatitis B

