



Influenza Vaccine Order Form 2009-2010 Season

VFC PIN

Fax order to (801) 538-9322

Date Submitted		Clinic Name			Phone with Area Code			
Delivery Address (no PO Boxes)							<input type="checkbox"/> Check if new address	
Person Completing Order (Print)				VFC Contact Person (Print)				<input type="checkbox"/> Check if new VFC Contact
All sections must be completed for your order to be processed. Orders submitted without complete VFC inventories will not be accepted.								
VACCINE	ALL VFC DOSES IN INVENTORY	BRAND NAME (MANUFACTURER)	DOSES ORDERED	PACKAGING	VFC PROGRAM USE ONLY			
					Doses Filled	Doses Back Ordered	VacMan Entry Date	
ALL VACCINES STORED IN THE REFRIGERATOR AT 35° - 46° F (2° - 8° C)								
Influenza Preservative Free Injectable		Fluzone-PF (Sanofi Pasteur) <i>*Ages 6-35 months only</i>		0.25 mL single dose syringes - 10 per box				
		Fluzone-PF (Sanofi Pasteur) <i>*Ages 36 months and older</i>		0.5 mL single dose vial - 10 per box				
Influenza Injectable		Fluzone (Sanofi Pasteur) <i>*Ages 6 months and older</i>		5 mL 10-dose vial - 1 vial per box				
		Fluvirin (Novartis) <i>*Ages 4 years and older</i>		5 mL 10-dose vial - 1 vial per box				
		Fluarix (GSK) <i>*18 years and older</i>		0.5 mL single dose syringes - 5 per box				
Influenza Preservative Free Intranasal		FluMist (MedImmune) <i>*Healthy persons 2-49 years</i>		0.2 mL single-use sprayers - 10 per box				
**Note: If the vaccine brand selected is not available, orders will be filled with a vaccine brand in inventory.								

Form 3C 08/09

Reminder on the ACIP Recommendations for Influenza

- ◆ Children 6 months through 8 years of age who received only 1 dose in their first year of vaccination should receive 2 DOSES the SECOND YEAR they are vaccinated.
- ◆ Children 6 months through 8 years who are in the third year or more of being vaccinated and who received only 1 dose in each of their first 2 years of being vaccinated should continue receiving a single annual dose.
- ◆ Two doses of Trivalent Inactivated Influenza Vaccine (TIV) should be separated by at least 4 weeks.
- ◆ Two doses of Live Attenuated Influenza Vaccine (LAIV) should be separated by at least 4 weeks.

Storage of All Influenza Vaccines

All Influenza vaccines, including FluMist, should be stored REFRIGERATED, not frozen, at 2 - 8° C (35 - 46° F) upon receipt and remain at that temperature until the expiration date.

For more information, please refer to the MMWR July 31, 2009, Prevention and Control of Seasonal Influenza with Vaccines; the product's package insert; contact the vaccine manufacturer; or call the Utah Immunization Program at (801) 538-9450.